

# **Analyst and investor fibrosis briefing**

A new class of protein therapeutics to treat human disease

February 2017

### **Analyst & investor fibrosis briefing**

- AdAlta's inaugural investor and analyst briefing aims to educate analysts and investors on our R&D technology platform as it relates to the science and the commercial opportunity to deliver value to shareholders
- ▶ AD-114 for idiopathic pulmonary fibrosis remains a core priority for the business
- AdAlta's focus is the completion of Phase I clinical studies and securing a global licensing deal for our lead i-body candidate
- Additional applications for the i-body platform in a range of other fibrotic diseases will also be discussed in detail

### **Analyst & investor fibrosis briefing**

- Recent international partnerships and collaborations have independently assessed and validated our technology
- Expert presentation by non-executive Director Dr Robert Peach will outline what it takes to get an antibody to Phase III and an \$8B acquisition
- AdAlta Chief Scientific Officer Mick Foley will outline AD-114 data in various therapeutic areas of fibrosis
- Presentations by leading researchers in lung, kidney and eye fibrosis will provide an overview of the clinical and commercial opportunities in this therapeutic area
- AdAlta's world class SAB John Westwick and Brian Richardson and AdAlta's non-executive Director Robert Peach will be available to answer specific questions and to discuss what it takes to get a fibrosis drug to the clinic and a deal



### **Agenda**



Robert Peach From NZ to the US, a Phase III antibody and an \$8B acquisition

Robert will discuss the Receptos story and what it takes to get an \$8B acquisition



Mick Foley

AD-114 a novel i-body for the treatment of fibrosis

Mick will provide a summary of AD-114 data in therapeutic areas of lung, liver and eye fibrosis



Glen Westall Pulmonary fibrosis – current state of play in Idiopathic Pulmonary Fibrosis

Glen will discuss what is IPF, how it is diagnosed, what the current treatments and their limitations and the drug development pipeline



Muh Geot Wong Renal fibrosis and chronic kidney disease

Muh Geot will discuss what happens with chronic kidney disease, how it is diagnosed, what the current treatments and their limitations and the drug development pipeline



### **Agenda**





Eye Fibrosis, causes, diseases and treatments

Erica will discuss eye fibrosis and the various diseases including wet-AMD, what the current treatments and the drug development pipeline



# PANEL DISCUSSION: Brian Richardson John Westwick Robert Peach

#### Getting a fibrosis drug to the clinic and a deal

The panel includes a number of drug development experts who have significant experience taking a drug from the research bench, through the clinic and providing to patients.

The panel will discuss what it takes to get a fibrosis drug to the clinic.

They will also discuss what it takes to get a deal with a pharmaceutical company.





#### AdAlta Investor Update

Sam will provide investors an update on AdAlta's achievements since the oversubscribed IPO in August and upcoming milestones







# Receptos story – From NZ to the US, a Phase III antibody and an \$8B acquisition

Robert Peach



### **Dr Robert Peach**

- ▶ Dr Peach has over 25 years of drug discovery and development experience in the pharmaceutical and biotechnology industry. In 2009 he co-founded Receptos, becoming Chief Scientific Officer and raising \$59M in venture capital and \$800M in an IPO and three subsequent follow-on offerings.
- ▶ In August 2015 Receptos was acquired by Celgene for \$7.8B. His extensive drug discovery and development experience in autoimmune and inflammatory diseases, and cancer has resulted in multiple drugs entering clinical trials and 3 registered drugs. He currently serves on the Board of Directors of Innate Immunotherapeutics and Avalia Immunotherapies and is a consultant for several other biotechnology companies.
- ▶ Robert is the co-author of 70 scientific publications and book chapters, and 17 patents. He was educated at the University of Canterbury and the University of Otago, New Zealand.



Groundbreaking Science to the Clinic

### **Presentation Outline**

- Receptos experience
  - Starting a biotechnology company in San Diego
    - Licensing assets
    - Raising venture capital
    - Hiring staff and building a productive culture
  - Ozanimod: a small molecule drug for treating relapsing multiple sclerosis and inflammatory bowel diseases
    - Clinical data
    - Market analyses
  - Strategies that lead to an \$8B acquisition



### The Path Towards Receptos – Key Roles

- → 1991-2000 Discovery and development of Orencia and Nulojix at Bristol-Myers Squibb; Seattle and Princeton
- ◆ 2000-2003 Director Antibody Discovery at IDEC. Rituxan (anti-CD20) for autoimmune diseases
- 2003-2007 Senior Director, Oncology Discovery at merged Biogen Idec.
   Multiple programs advanced into clinic. In-licensing,
   acquisition, VC investments
- ◆ 2007-2009 Co-founder and VP Biology at Apoptos. Oncology assets licensed from Burnham Institute
- ◆ 2009-2015 Co-founder and Chief Scientific Officer of Receptos

### Receptos Start-Up: 2009

- Starting with a failure: Apoptos
- Receptos was founded in 2009 with technology licenses from The Scripps Research Institute™
- Focus primarily on autoimmune disease drug discovery and development, and a structure-based drug design technology platform
- → 2009: A tough time raising \$25M in venture capital
  - Venrock, Arch, Lilly, Flagship
- → \$34M in 2010
  - Polaris, Venrock, Arch, Lilly, Flagship
- → 20 scientists, 1 business development, 1 accountant, 1 admin
  - → Deep experience, multiple skill sets, cultural "fit"
  - No CEO, CMO, CFO etc



### Receptos Location: Torrey Pines Mesa in San Diego



### What did we License from Scripps?

- GPCR structural biology technology platform
  - Established industry partnerships that raised \$20M in non-dilutive capital
- Small molecule compound "hits" identified in a high through-put screen targeting the activation of a protein called sphingosine 1-phosphate 1 receptor (S1P1R)
- Receptor is expressed on lymphocytes and is involved in controlling the trafficking of these cells out of lymphoid tissues
- Biological rationale suggested an ability to impact pathology associated with multiple autoimmune diseases, including multiple sclerosis and inflammatory bowel diseases (ulcerative colitis, Crohn's disease)
- Clinical proof of concept established for this mechanism of action



## Drug Discovery Goals: Best in Class and Rapid Development


- Develop an oral daily pill that could be dosed chronically and had a superior safety profile to "same class" multiple competitor drugs, including Gilenya, that was close to completing clinical development
  - Cardiac toxicity including 1st dose heart rate monitoring
  - Prolonged drug activity
  - Induction of fibrosis
- Rapid preclinical development
  - 1 year to identify clinical candidate
  - → 10 months to file an IND (November 2010)
  - Phase 1 SAD/MAD began 1 month after IND approved (January 2011)

### RPC1063 (Ozanimod)


 Optimal potency, selectivity, oral bioavailability, absorption, distribution, metabolism, excretion, toxicity, pharmacokinetic and pharmacodynamic properties

### **RADIANCE Phase 2/3 Trial Overview**


### Relapsing Multiple Sclerosis SPA Agreement with FDA



### Phase 2 Primary Endpoint: Mean Cumulative Number of GdE Lesions From Wk 12 to Wk 24



# First Dose Effect on Mean Heart Rate Hours 1 to 6: Comparison to Gilenya\*



\*Data estimated from Novartis material at MSBoston2014; not a head to head comparison


### Current Clinical Trial Status in Relapsing Multiple Sclerosis

- Two large phase 3 registrational trials ongoing
  - Worldwide
  - → 1,300 patients per trial
  - 1 year of dosing
  - 2 years of dosing
  - Compared to Avonex, a currently approved drug
  - \$100,000/patient; \$130M/trial
  - Clinical development completed in 2017
  - New Drug Application and approval with FDA/Europe in 2018




## RPC1063 is Active in a Therapeutic SCID Mouse Model of Inflammatory Bowel Disease

CD4+ CD45RBhigh T-Cell Adoptive Transfer: Histopathology



# Phase 2 Ulcerative Colitis: Proportion of Patients in Remission at Week 8 (April 2015)


Primary Endpoint




### The Relapsing Multiple Sclerosis Market is Valued at ~\$16B

**UNMET NEEDS** 








More tolerable & convenient therapies

Agents that halt or reverse damage


### Oral Share of RMS IMS Scripts Rapidly Increasing Gilenya® and Tecfidera® drive rapid increases in oral market share gains



Note: RMS Therapies Include Avonex®, Betaseron®, Copaxone®, Rebif®, Gilenya®, Aubagio® & Tecfidera®; Orals = Gilenya®, Aubagio® & Tecfidera®; ABCRs = Avonex®, Betaseron®, Copaxone® & Rebif®




### RPC1063 Poised to be Next S1P<sub>1.5</sub>R Modulator into RMS Market




### RMS Market Valued at \$16B and Growing Marketing Strategy for RMS Includes Competing Head to Head with Tecfidera® in Front Line While "Beating" Gilenya® in Later Lines


### **Positioning of MS Treatment Options**



## In Market Research, RPC1063 Profile Viewed More Favorably Than Tecfidera® on Efficacy and Tolerability and Gilenya® on Safety



### RPC1063 Viewed More Favorably Than Gilenya® on Safety




- Majority of physicians had a positive view of RPC1063 on all metrics tested compared to currently marketed orals
- When shown patient profiles, physicians selected RPC1063 more frequently than any marketed RMS therapy

Note: For purposes of the survey, respondents were provided with a product profile for RPC1063 that assumed efficacy equal to Gilenya® with specified improvements in safety profile; Actual clinical results for late stage trial and the FDA marketing label for RPC1063, if approved, may be different than the profile that was assumed for market research; Pricing and reimbursement were assumed to be equal among RPC1063 and all other RMS treatments

# The UC Market Opportunity for RPC1063 in the US: Gastroenterologists Position RPC1063 in Early Lines Against Immunosuppressants *as well as* Last Line Against Anti-TNFs

#### **Positioning of UC Treatment Options in Market Research Outcomes**



Positioning based on Receptos sponsored market research. Respondents were provided with a product profile for RPC1063 and vedolizumab that assumed comparable efficacy improved to anti-TNFs with specified improvements in safety profile. Actual clinical results for late stage trial and the FDA marketing label for RPC1063, if approved, and vedolizumab, approved in May 2014, may be different than the profile that was assumed for market research; pricing and reimbursement were assumed to be equal among RPC1063 and all other UC treatments Source: Kantar 2013, US Gastroenterologist Quantitative Survey (N = 101)

## Market Research Results: US Gastroenterologist Estimated RPC1063 Patient Share in IBD

Ulcerative Colitis		Treatment Algorithm	Crohn's D	Crohn's Disease	
RPC1063 Patient Share	Current Treatment Choice by Line		Current Treatment <a href="Choice by Line">Choice by Line</a>	RPC1063 <u>Patient Share</u>	
7% *	5ASA →	First Line	◆5ASA and	7.0% *	
		•	Corticosteroids		
13.0%	Corticosteroids 🔸	Second Line	→6-MP, AZA	14.0%	
		•			
18.0%	6-MP, AZA →	Third Line	<b>→</b> Biologics	20.0%	
		•			
24.0%	Biologics 🔸	Last / Fourth Line	<b>→</b> Biologics	27.0%	

Note: \* Physicians were asked to estimate first line RPC1063 peak patient share IF clinical trial data was available demonstrating improvement to 5-ASAs. Receptos applies appropriate discounts to market research results for the purposes of financial modeling.



### Receptos: 2015

- → IPO (NASDAQ: RCPT) in May 2013. \$83M raised. \$14/share
- → Additional 3 capital raises in 2014, totaling ~\$700M. Each raise over-subscribed
- 110 employees and growing
- → Phase 3 clinical trials in relapsing multiple sclerosis and phase 2 in inflammatory bowel diseases (UC and Crohn's disease)
- ◆ Phase 2 in EoE
- Preclinical program in T2D/NASH
- Discovery chemokine receptor program
- Acquired July 2015 for \$7.8B
  - \$232/share



### **Key Strategies and Lessons Learnt**

- Hire the "right" people and on an "as needed" basis. Wear lots of hats and preserve \$\$
- Build a productive, creative, rewarding culture
- Focus, don't waste time or money. Understand the critical path activities
- Do rigorous science
- Don't partner early
- Enact bold (but not reckless) clinical development plans
- Understand the competition and where your drug might fit in the market place
- Under promise and over deliver
- Build relationships: FDA, CRO's, consultants, KOL's, investors, pharma/biotech
- Raise money when you can
- Maintain optionality



### Kaikoura





### AD-114 a novel i-body for the treatment of fibrosis

Mick Foley, CSO AdAlta Limited m.foley@adalta.com.au



### A/Prof Mick Foley

- Mick is the founding scientist of AdAlta and a key inventor of AdAlta's lead i-body candidate AD-114.
- ▶ Upon completion of his PhD he was awarded a Wellcome Training Fellowship and worked at the Walter and Elisa Hall Institute. In 1995 Mick was awarded an ARC QEII Fellowship where he established the phage display of antibodies and peptide technology as a means of answering fundamental questions of immunity to infectious diseases.
- Mick is an internationally recognized leader in phage display, the technology used to screen the i-body library to identify new drug candidates. Having published over 70 scientific publications Mick has received funding from ARC, NHMRC and NIH (US).

### **Disclaimer**

Investment in AdAlta is subject to investment risk, including possible loss of income and capital invested. AdAlta does not guarantee any particular rate of return or performance, nor do they guarantee the repayment of capital.

This presentation is not an offer or invitation for subscription or purchase of or a recommendation of securities. It does not take into account the investment objectives, financial situation and particular needs of the investor. Before making any investment in AdAlta, the investor or prospective investor should consider whether such an investment is appropriate to their particular investment needs, objectives and financial circumstances and consult an investment advisor if necessary.


This presentation may contain forward-looking statements regarding the potential of the Company's projects and interests and the development and therapeutic potential of the company's research and development. Any statement describing a goal, expectation, intention or belief of the company is a forward-looking statement and should be considered an at-risk statement. Such statements are subject to certain risks and uncertainties, particularly those inherent in the process of discovering, developing and commercialising drugs that are safe and effective for use as human therapeutics and the financing of such activities. There is no guarantee that the Company's research and development projects and interests (where applicable) will receive regulatory approvals or prove to be commercially successful in the future. Actual results of further research could differ from those projected or detailed in this presentation. As a result, you are cautioned not to rely on forward-looking statements. Consideration should be given to these and other risks concerning research and development programs referred to in this presentation.



# Fibrosis: unmet medical need with multiple indications

- Developing i-bodies as improved therapies for the treatment of fibrosis
  - a condition that is prevalent in 45-50% of all diseases
- Fibrosis can occur in many tissues of the body as a result of inflammation or damage
  - it can result in scarring of vital organs causing irreparable damage and eventual organ failure
- AdAlta's initial focus is on lung fibrosis

Collectively fibrosis represents a large unmet clinical need



### i-body technology


AdAlta is developing a new technology platform that produces unique proteins known as i-bodies, that mimic the shape of shark antibody binding domain and engineers their key stability features into a human protein, for therapeutic intervention in disease.

The single domain antigen binding region of shark antibodies is extremely stable and has a long binding loop not present in either human or next generation antibodies.

#### Advantages of i-bodies

- High target specificity and high affinity for their target
- Small proteins; 10% the size of a typical human antibody
- Highly stable to proteases, high temperatures and low pH
- Long loop that can bind to a diverse range of therapeutically relevant targets including those that are difficult for current antibody therapies
- ▶ <u>Human protein</u> reduced risk of immune response

Long loop that enables access to novel drug targets



**Shark Antibody** 



# AD-114 lead program in Idiopathic Pulmonary Fibrosis (IPF)

- ► AD-114 is lead i-body candidate in pre-clinical development
  - Demonstrates both anti-fibrotic and anti-inflammatory activity in the lung
  - Important for arresting and modifying the disease and tackling the treatment of idiopathic pulmonary fibrosis (IPF); this is the primary indication



**IPF** 

#### **Idiopathic Pulmonary Fibrosis**

A chronic, highly lethal and rare disease. 50-70% mortality rate >135,000 people in US alone World wide sales ~\$4.2B by 2020

Source: Evaluate Pharma, Orphan Drug Report 2015



# CXCR4 and idiopathic pulmonary fibrosis (IPF)


Patients that rapidly progress express more CXCR4 compared to slow IPF progressors

CXCR4 +ve cells (fibrocytes) significantly elevated in stable IPF patients, and further increased during acute exacerbations


Fibrocytes not correlated with lung function but an independent predictor of early mortality

- ▶ 7.5 months with more than 5% fibrocytes
- ▶ 27 months with less than 5% fibrocytes

#### CXCR4 expression increased in fast progressing IPF patient tissue



#### Fibrocyte numbers predict mortality




REF: Moeller, et al. Am J Respir Crit Care Med Vol 179. pp 588–594, 2009




## AD-114 binds to lung tissue from patients with fibrosis

AD-114 was used for Immunohistochemical (IHC) staining of normal and diseased lung tissues to verify expression of CXCR4 *in situ* 



AD-114 does not bind lung tissue from normal lungs



AD-114 binds to lung tissue from lungs with fibrosis





## Migration/invasion specifically reduced with IPF lung fibroblasts


AD-114 specifically inhibited migration of slow and rapid IPF fibroblast migration but did not have any effect on normal fibroblasts.

AD-114 has greater *in vitro* efficacy compared to the only approved therapies Nintedanib and Pirfenidone for IPF treatment.


MIGRATION	No effect on normal fibroblasts	Inhibits slow IPF progressors	Inhibits fast IPF progressors
i-body AD-114	<b>V</b>	<b>V</b>	<b>✓</b>
Nintedanib (Boehringer)	X	<b>V</b>	<b>V</b>
Pirfenidone (Roche)	<b>/</b>	X	X
Other CXCR4 drug (Sanofi)	<b>V</b>	X	X









**Prophylactic or Preventative setting** 



Normal lung tissue



**IPF lung tissue** (lung disease mouse model)




IPF lung tissue + AD-114 dosed for 21 days (lung disease mouse model)


AD-114 reduces collagen content and inflammatory cell infiltration in the Bleomycin mouse model and demonstrates a similar architecture to that of the normal lung

**Prophylactic or Preventative setting** 

► AD-114 significantly reduced the Ashcroft score compared to the Bleomycin treated mice


The negative i-body at the same dose as AD-114 had no effect on preventing fibrosis









Therapeutic setting



Normal lung tissue



IPF lung tissue (lung disease mouse model)




IPF lung tissue + AD-114 dosed for 13 days (dosing started at day 8 in lung disease model)

AD-114 reduces collagen content and inflammatory cell infiltration in the Bleomycin mouse model and demonstrates a similar architecture to that of the normal lung




#### Therapeutic setting

- ► AD-114 significantly reduced the Ashcroft score compared to the Bleomycin treated mice
- AMD3100 (CXCR4 antagonist) also reduced the the Ashcroft score compared to the Bleomycin treated mice



#### **Ashcroft Score Therapeutic Mode**



# AD-114 and non-alcoholic steatohepatitis (NASH)

- Non-alcoholic steatohepatitis (NASH) is a pandemic, metabolic disease which has both inflammatory and fibrotic components
- ► AD-114 is lead i-body candidate in pre-clinical development
  - Demonstrates both anti-fibrotic and anti-inflammatory activity in the liver
  - Important for arresting and modifying the disease and tackling the treatment of NASH




Liver
NASH & CIRRHOSIS

#### **NASH**


A chronic disease with high levels of morbidity and mortality About 3-5% of adults in the United States have NASH Sales of drugs for the treatment of fibrosis caused by NASH are estimated to be US\$1.6 billion by 2020.

## AD-114 prevents fibrosis in a mouse model of liver fibrosis


Therapeutic setting



Normal liver tissue



NASH liver tissue (liver disease mouse model)




NASH liver tissue + AD-114 dosed for 21 days (liver disease mouse model)

AD-114 significantly reduces hepatocellular ballooning, a key feature required for the diagnosis of NASH



## AD-114 prevents fibrosis in a mouse model of liver fibrosis

- AD-114 decreased serum ALT levels and non-alcoholic fatty liver disease (NAFLD) score compared with the vehicle or disease model group
- ► The improvement in serum ALT levels suggests that i-body ameliorated hepatocellular injury and inflammation preventing progression of disease
- Hepatocyte ballooning was significantly decreased compared with the vehicle or diseased group
- AD-114 possess hepatoprotective and anti-NASH effects





### **AD-114** and eye fibrosis

- Infections or inflammation in the eye result in impairment of visual function and can ultimately lead to fibrosis. Complications from common eye diseases that can result in fibrosis occur in age related macular degeneration (AMD) and diabetic retinopathy.
- ▶ AD-114 is lead i-body candidate in pre-clinical development
  - Demonstrates both anti-fibrotic and anti-leakage activity in the eye
  - Important for arresting and modifying the disease and tackling the treatment of eye fibrosis



Eye Wet-AMD & PVR

#### **Eye Fibrosis**


AMD is the commonest cause of severe visual impairment in people over the age of 50 years in the developed world >1m in AU and 2m in USA with AMD Market research estimates that the market size for AMD will be over US\$10 billion by 2023 while the market size for diabetic retinopathy will be US\$10 billion in 2022.




### **AD-114** prevents eye fibrosis


- Mouse chorodial neovascularization model(CNV): laser burn to the retina
  - Induces subretinal haemorrhage
  - Contraction of retinal tissue
  - Alteration in microglia and glial response
  - Alteration in gene expression
- IVT injection of single dose of ibody
  - Improves retinal retraction and reduces lesion size
  - Fibrosis gene expression reduced

#### AD-114 reduces contraction and lesion size in eye fibrosis mouse model






#### No Treatment



**Treatment with AD-114** 










#### AD-114 reduces lesion size and number

- ▶ AD-114 is able to reduce the number and size of the lesion in both preventative and therapeutic models of wet-AMD (CNV)
- ▶ AD-114 is able to significantly reduce fibrosis as measured by trichrome staining in both preventative and therapeutic models of wet-AMD (CNV)



#### AdAlta summary

- Powerful proprietary technology platform to develop a pipeline of i-bodies for the treatment of a wide range of human diseases
  - Extreme stability of i-body similar to single domain shark antibody
  - Long loop of i-body binds deep in GPCR pocket and has functional activity
- Advanced lead candidate AD-114 with significant pre-clinical validation
  - has specificity for diseased human tissue with effects only shown on IPF tissue and no effects displayed on normal lung tissue nor any evidence of off target effects;
  - is more effective than existing IPF approved drugs showing greater in vitro efficacy compared to the only approved therapies Nintedanib and Pirfenidone;
  - demonstrates both anti-fibrotic and anti-inflammatory effects in multiple animal models in multiple areas of fibrosis; and
  - is a novel mechanism of action for fibrosis making AD-114 a potential "first in class" therapy.





## Pulmonary fibrosis – current state of play in Idiopathic Pulmonary Fibrosis

#### A/Prof Glen Westall

Lung Fibrosis Service, Alfred Hospital

NHMRC Centre of Research Excellence: Lung Fibrosis



#### A/Prof Glen Westall


- ▶ Dr Glen Westall received his undergraduate medical training at King's College Medical School in London, UK, before training in general and respiratory medicine at the Royal Brompton Hospital in London and the Alfred Hospital in Melbourne. His clinical interests include advanced lung disease, bronchoscopy and lung transplantation. He is physician-in-charge of the Paediatric Lung Transplant program at the Alfred.
- ▶ Glen's research interests parallel his clinical expertise in advanced lung disease and lung transplantation. He has subsequently developed a wider interest into how activation of the innate immune system early post-lung transplant influences clinical outcomes. Establishing a research platform in pulmonary xenotransplantation has resulted in the award of an internationally competitive Career Development Award. Clinically, Glen is a principal and co-investigator on numerous studies in lung transplantation and bronchoscopic lung volume reduction.

### **Lung Fibrosis**


## Scaring of the Lung

Lungs become stiff

Lungs shrink in size




**Normal CXR** 



**Lung Fibrosis** 

### Interstitial lung disease (ILD) classification





### **Idiopathic Pulmonary Fibrosis (IPF)**



Represents 50% of all Fibrotic Lung Conditions

Male:Female 1.5:1

Age of Presentation 66 yrs

Smoking history 70%

Australian IPF Cases 5000-10,000

Diagnosed by MDT Meeting




IPF Diagnosis: Multi-Disciplinary Team (MDT)

meeting

 Communication between clinician, radiologist and when appropriate pathologist

- Clinical data
  - Presentation, Exposures, Smoking status, Associated disease, Lung function and Radiologic findings




THE



### Traditional Therapy for IPF: historical view

- Uncontrolled Chronic Inflammation
  - Corticosteroids
  - Cyclophosphamide
  - Azathioprine
  - N-acetylcysteine (NAC)



# 2011 Evidence Based Guidelines: Treatment recommendations

Agent	For	Against
Corticosteroids		Strong +
Colchicine		Strong +
Cyclosporine A		Strong +
Corticosteroid and imm-mod		Strong ++
Interferon gamma		Strong ++++
Bosentan		Strong +++
Etanercept		Strong +++
NAC, predisolone, azathioprine		Weak ++
NAC monotherapy		Weak ++
Anticoagulation		Weak +
Pirfenidone		Weak ++

Raghu et al. Am J Respir Crit Care Med Vol 183. pp 788-824, 2011



## 2011 Evidence Based Guidelines: Treatment recommendations

Agent	For	Against
Corticosteroids		Strong +
Colchicine		Strong +
Cyclosporine A		Strong +
Corticosteroid and imm-mod		Strong ++
Interferon gamma		Strong ++++
Bosentan		Strong +++
Etanercept		Strong +++
NAC, predisolone, azathioprine		Weak ++
NAC monotherapy		Weak ++
Anticoagulation		Weak +
Pirfenidone		Weak ++



Raghu et al. Am J Respir Crit Care Med Vol 183. pp 788-824, 2011


### 2014: A big year for lung fibrosis!

## Pirfenidone approved by FDA





#### Reduction in mortality and decreased FVC



47% reduction in proportion of patients with a decline in FVC

NEJM 2014; 370: 2083

Pooled data: Ascend & Capacity studies

All cause mortality 

48%

IPF deaths 

68%

#### **Adverse Events**

Adverse Event	Pirfenidone (N = 278)	Placebo (N = 277)	
	no. of patients (%)		
Cough	70 (25.2)	82 (29.6)	
Nausea	100 (36.0)	37 (13.4)	
Headache	72 (25.9)	64 (23.1)	
Diarrhea	62 (22.3)	60 (21.7)	
Upper respiratory tract infection	61 (21.9)	56 (20.2)	
Fatigue	58 (20.9)	48 (17.3)	
Rash	78 (28.1)	24 (8.7)	
Dyspnea	41 (14.7)	49 (17.7)	
Dizziness	49 (17.6)	36 (13.0)	
Idiopathic pulmonary fibrosis†	26 (9.4)	50 (18.1)	
Bronchitis	39 (14.0)	36 (13.0)	
Constipation	32 (11.5)	38 (13.7)	
Back pain	30 (10.8)	37 (13.4)	
Dyspepsia	49 (17.6)	17 (6.1)	
Nasopharyngitis	33 (11.9)	30 (10.8)	
Anorexia	44 (15.8)	18 (6.5)	
Vomiting	36 (12.9)	24 (8.7)	
Decrease in weight	35 (12.6)	22 (7.9)	
Gastroesophageal reflux	33 (11.9)	18 (6.5)	
Insomnia	31 (11.2)	18 (6.5)	



### 2014: A big year for lung fibrosis!

#### Nintedanib approved by FDA




#### Reduction in mortality and decreased FVC

Intracellular inhibitor of multiple tyrosine kinases

#### **NB: Not curative**

Secondary end points not achieved



#### **Adverse Events**


means reconstruction of the control	INPUL	ere s	INPUL	ere s	
Event					
	Nintedanib (N = 309)	Placebo (N = 204)	(N = 329)	Placebo (N=219)	
	number of patients (percent)				
Any adverse event	298 (96.4)	181 (88.7)	311 (94.5)	198 (90.4	
Any adverse event, excluding progression of idiopathic pulmonary fibrosis*	296 (95.8)	179 (87.7)	311 (94.5)	197 (90.0	
Most frequent adverse events†					
Diarrhea	190 (61.5)	38 (18.6)	208 (63.2)	40 (18.3	
Nausea	70 (22.7)	12 (5.9)	86 (26.1)	16 (7.3)	
Nasopharyngitis	39 (12.6)	34 (16.7)	48 (14.6)	34 (15.5	
Cough	47 (15.2)	26 (12.7)	38 (11.6)	31 (14.2	
Progression of idiopathic pulmonary fibrosis*	31 (10.0)	21 (10.3)	33 (10.0)	40 (18.3	
Bronchitis	36 (11.7)	28 (13.7)	31 (9.4)	17 (7.8)	
Upper respiratory tract infection	28 (9.1)	18 (8.8)	30 (9.1)	24 (11.0	
Dyspnea	22 (7.1)	23 (11.3)	27 (8.2)	25 (11.4	
Decreased appetite	26 (8.4)	14 (6.9)	42 (12.8)	10 (4.6)	
Vomiting	40 (12.9)	4 (2.0)	34 (10.3)	7 (3.2)	
Weight loss	25 (8.1)	13 (6.4)	37 (11.2)	2 (0.9)	
Severe adverse events‡	81 (26.2)	37 (18.1)	93 (28.3)	62 (28.3	
Serious adverse events‡	96 (31.1)	55 (27.0)	98 (29.8)	72 (32.9	
Fatal adverse events	12 (3.9)	10 (4.9)	25 (7.6)	21 (9.6)	
Adverse events leading to treatment discontinuation§	65 (21.0)	22 (10.8)	58 (17.6)	33 (15.1	
Gastrointestinal disorders	26 (8.4)	3 (1.5)	21 (6.4)	2 (0.9)	
Respiratory, thoracic, and mediastinal disorders	12 (3.9)	10 (4.9)	8 (2.4)	18 (8.2)	
Investigation results¶	10 (3.2)	1 (0.5)	8 (2.4)	1 (0.5)	
Cardiac disorders	5 (1.6)	4 (2.0)	2 (0.6)	3 (1.4)	
General disorders and conditions involving site of study-drug administration	8 (2.6)	3 (1.5)	2 (0.6)	1 (0.5)	

Study discontinuation 5%



### Unmet clinical need: drugs in development

- Current drugs have limited efficacy and substantial side effects
- Other anti-fibrotic and anti-angiogenic agents currently in development with multiple targets
  - Inhaled Pirfenidone
  - WNT signaling
  - FG-3019
  - Zileutin
  - ACE inhibitors
  - Anti-vβ6 integrin
  - Anti-IL13
  - Anti-IL13/IL4
  - Imatanib
  - LOXL-2





### Clinical trials.gov

Phase 1 Dasatinib (oral Bcr-Abl TK inhibitor, BMS) + Quercetin

Vismodegib (hedgehog signalling pathway) + perfinidone (Genentech, Roche)

CC-90001 (Celgene, JNK inhibitor)

Mesenchymal stem cells

Phase 2 Rituximab

Tipelukast (medicnova)

STX-100 ( $\alpha v \beta 6$  integrin inhibitor, Biogen)

GBT440 (oxygen carrying capacity, Global blood therapeutics)

KD0125 (ROCK2 signalling inhibitor, Kardmon)

Lebrikizumab and perfinidone (Roche)

GLPG1690 (autotaxin inhibitor)

Phase 3 Bactrim

Cyclophosphamide (acute exacerbations)

Nintedanib and sildenafil (BI)


Perfinidone and sildenafil (Roche)

ART-123 (acute exacerbations) thrombomodilin

Phase 4 perfinidone + nintedanib



### **Drug Discovery and translational research**



Assessment of human IPF tissue provides additional evidence of efficacy prior to clinical trials

#### **Alfred Tissue Bank**



- National resource
- Collaborations (Academic and Commercial entity)
- Creation of pilot data
- ► Conduit: Alfred Lung Transplant Program (n=90)




ILD Explant Lungs



Normal Lungs donated for research (Donate Life)

#### **CXCR4** Role in IPF

Very limited expression in normal or nondiseased tissues







Figure S. Immunohistochemical staining of IPF lung tissue for CXCR4 (brown). 8x magnification, scale bars are 500 µm.

CXCR4 is upregulated in IPF tissue



### Summary

- Increasing understanding of the pathophysiology
- Treatments (albeit limited)
- Many Lung Fibrosis services/clinics



Need better therapies for a terminal disease!!





#### Renal fibrosis and chronic kidney disease

Novel therapeutic targets for kidney fibrosis

#### **Professor Muh Geot Wong**

Royal North Shore Hospital, University of Sydney Australia George Institute for Global Health



### Dr Muh Geot Wong

- ▶ Dr Wong is Renal Physician at the Royal North Shore hospital, Sydney, Australia. He is also a senior research fellow at the George Institute of Global Health and a senior clinical lecturer at the University of Sydney. His PhD entitled "Novel therapeutic options in models of nephropathy" was awarded in 2011.
- ► His main area of research is in understanding the pathomechanisms of kidney tubulointerstitial fibrosis and biomarkers in predicting progression of chronic kidney disease particularly in diabetic kidney disease.

#### DISCLOSURES

- Speaker fees; Astra Zeneca
- The George Institute for Global Health holds research contracts for trials in cardiovascular and/or kidney disease with several organization including Boehringer Ingelheim, Merck, AbbVie, Roche, AstraZeneca, Servier, Astellas, Baxter, BMS, GSK, Janssen and Pfizer



#### **Overview**

- Residual risk of standard of care
- Mechanism of kidney fibrosis
- Novel therapies for kidney fibrosis
- Global market potential









## **Kidney Fibrosis**

Kidney or renal fibrosis is seen in virtually all progressive kidney diseases including diabetic nephropathy, allograft nephropathy or aging



Normal Kidney




**CKD** 






### Causes of end stage kidney disease over time

#### **AUSTRALIA**



#### **USA**










#### Renin Angiotensin Aldosterone system











# Residual risk of end stage renal disease on conventional therapy
















## Chronic kidney disease associated pathways are shared between renal diseases



Martini S et al. JASN 2013



Drug Discovery Today. March 2015

#### **Novel therapies and status**

- Anti TGFB antibody (LY2382770): Negative Study
- Pirfenidone: Study withdrawn
- Nox1/4 inhibitor: negative trial
- ► Tranilast and analogues FT011: In Phase 1b Clinical Trial
- Anti-CTGF Antibodies FG3019: Studies terminated
- CCR2/MCP1 inhibition: Larger clinical trial with hard renal endpoints needed
- SSAO/ VAP1 inhibitors: Phase 1 Clinical Trial concluded. ? Results
- ► Tie2Rec activator: angiopoietin receptor, tyrosine kinase inhibitor in development
- JAK-STAT inhibitors: in development
- ▶ LOX inhibitors: in development



CXCR4 and AdAlta's i-body?






# CXCR4 promising therapeutic strategy for treatment of kidney fibrosis

- CXCR4 expression was increased in disease state compared with normal tissue
- Collaboration between Kolling Renal Research and AdAlta ibody technology in exploring CXCR4 as therapeutic strategy to prevent the development of fibrosis

#### CXCR4 in kidneys of diabetic patients









### Challenges for clinical trial in CKD/Fibrosis

- Patients with CKD usually excluded from major trial
- Lack of intermediate endpoints/ biomarkers that regulatory agencies will accept for approval
- High trial failure rate
- ► Lack of coordination between scientists, investors, clinical trialists, pharma companies, regulatory authorities, policy makers and governments to develop novel strategies.









### **EMPA-REG**


#### Primary outcome: 3-point MACE






#### Zinman et al. N Eng J Med Sept 17 2015

#### CV death



All-cause mortality







## Effects of empagliflozin on renal outcomes

Hazard ratio 95% Ci

<b>&gt;</b>	Incident/worsening of nephropathy	0.61	0.53 - 0.70	<0.001
<b>&gt;</b>	New onset macroalbuminuria	0.62	0.54 - 0.72	<0.0001
<b>&gt;</b>	Doubling creatinine or ESKD	0.54	0.40 - 0.75	0.0002
<b>&gt;</b>	Doubling of serum creatinine	0.56	0.39 - 0.79	0.0009
<b>&gt;</b>	ESKD	0.45	0.21 - 0.97	0.04

The impact of empagliflozin on the primary end-point was not diminished in patients with CKD compared to those without it (MACE HR 0.88, CV death HR 0.78, heart failure HR 0.59, all-cause mortality HR 0.80).









p value

#### SGLT2 inhibition: glycosuria and natriuresis













#### LEADER study: Liraglutide in type 2 diabetes






Number needed to treat (NNT) in 3 years to prevent primary outcome=66, and to prevent death from any cause=98

Marso et al NEJM July 2016





# Projecting the estimated number of patients receiving RRT from 2010 to 2030



### Global "push"

▶ International Society of Nephrology (ISN) established working groups to establish and validate novel therapeutic targets to retard progression of CKD.

#### ► Four goals

- Improve the identification of "druggable" targets that are amenable to therapeutics
- Enhance the capacity for pre-clinical and early clinical development
- Broaden the availability of novel therapeutic approaches
- Increase investment in the development of therapies to limit CKD







#### **Global Market Opportunity**

- According to Decision Resources Group, the CKD market will achieve total sales of \$11.7 billion in 2022.
- ▶ Erythropoietin-stimulating agents (ESAs), phosphate binders, calcium mimetics, active vitamin D analogues, antihypertensive agents, IV iron and emerging CKD therapies for the CKD non-dialysis and dialysis patient populations. This analysis takes into account the offset of increasing costs of new therapies by emerging availability of generic drugs
- ► The launch and uptake of novel drugs during the next ten years will be the largest driver of market growth.
- Of the emerging therapies with novel mechanisms of action, the hypoxia-inducible factor-prolyl hydroxylase (HIF-PH) inhibitors will likely have the biggest impact on the CKD market.

2012 Patient Population	
Late-stage CKD Population <sup>8</sup>	2,356,913
Treated Population <sup>b</sup>	1,446,904
2012 Market Sales	
US	\$1.48bn
5EU	\$397m
Total	\$1.88bn
Key events (2012–2017)	Level of
PA-21 launch in the US and EU - 2014	11
Zerenex launch in the US and EU – 2014/2015	11
Renagel/Renvela patent expiry in the US and EU – 2014	111
Oral treatments included in the Medicare dialysis reimbursement bundle – 2016	11
Velcalcetide launch in the US and EU – 2016/2017	11
2017 Market Sales	
US	\$1.27bn
5EU	\$391m
Total	\$1.66bn

a = Stage 4 and 5 CKD prevalence cases; b = patients treated for hyperphosphatemia and/or secondary hyperparathyroidism





#### **Summary**

- An unmet need for new therapeutic options of kidney fibrosis
- Global recognition, demand and commercial opportunity to develop new therapeutics in CKD
- Promising novel targets needs clinical translation and hard renal outcomes
- ➤ Collaborative effort among scientists, clinicians, investors, trialists, pharma companies, regulatory authorities, policy makers and governments needed









## Eye Fibrosis, causes, diseases and treatments

### Professor Erica L Fletcher

Department of Anatomy and Neuroscience The University of Melbourne


Email: elf@unimelb.edu.au




#### **Prof Erica Fletcher**

- ► Erica Fletcher is Professor in the Department of Anatomy and Neuroscience, at The University of Melbourne where she heads the Visual Neuroscience Laboratory.
- ▶ She is a clinically trained optometrist who holds both MSc and PhD degrees. She completed her PhD at The University of Melbourne and undertook postdoctoral training at the Max Planck Institute for Brain Research in Germany, funded by a CJ Martin Award from the NH&MRC.
- ▶ Prof Fletcher was appointed to an academic position in 2000 at The University of Melbourne. Prof Fletcher's research interests remain primarily focussed on understanding the causes of retinal disease.

## The eye












## Fibrosis and the eye

- Definition:
  - Scarring of the retina leading to vision loss
- ► Diseases of the retina associated with fibrosis:



- Diabetic retinopathy and Age related macular degeneration.
- -Retinal detachment:
  - Fibrosis (PVR): 8-10% of all patients undergoing primary retinal detachment surgery











#### The retina and its vasculature



{Image: Prof Erica Fletcher Univ Melb}






{Image: Friedlander, 2007 J Clin Invest}




## Blindness: 50-100,000 Australians




## Age related macular degeneration





{Image: Prof Robyn Guymer, CERA}





{Image: Prof Robyn Guymer, CERA}





{Image: Prof Robyn Guymer, CERA}

# First medical treatment for wet AMD 2002










{Image: Prof Robyn Guymer, CERA}

#### **Anti-VEGF treatments**







Brown et al, N Engl J Med 2006; 355(14); 1432

#### Incidence rates of legal blindness from AMD

Denmark population based observational registry study



## Long term effects: A sting in the tail!



## There is a problem: fibrosis

## Risk of Scar in the Comparison of Age-related Macular Degeneration Treatments Trials

Ebenezer Daniel, MBBS, PhD, <sup>1</sup> Cynthia A. Toth, MD, <sup>2</sup> Juan E. Grunwald, MD, <sup>1</sup> Glenn J. Jaffe, MD, <sup>2</sup> Daniel F. Martin, MD, <sup>3</sup> Stuart L. Fine, MD, <sup>4</sup> Jiayan Huang, MS, <sup>1</sup> Gui-shuang Ying, MD, PhD, <sup>1</sup> Stephanie A. Hagstrom, PhD, <sup>3</sup> Katrina Winter, BS, <sup>2</sup> Maureen G. Maguire, PhD, <sup>1</sup> for the Comparison of Age-related Macular Degeneration Treatments Trials Research Group\*


### Results: Scar developed in 480 of 1059 eyes (45.3%) by 2 years.

versus occult CNV, blocked fluorescence (aHR, 1.4; CI, 1.1–1.8), toveal retinal thickness >212 μm (aHR, 2.4; CI, 1.7–3.6) versus <120 μm, foveal subretinal tissue complex thickness >275 μm (aHR, 2.4; CI, 1.7–3.6) versus ≤75 μm, foveal subretinal fluid (aHR, 1.5; CI, 1.1–2.0) versus no subretinal fluid, and subretinal hyperreflective material (SHRM) (aHR, 1.7; CI, 1.3–2.3) versus no SHRM. Eyes with elevation of the retinal pigment epithelium







### Reducing vision in wet AMD



Exudative (wet) AMD





Scarring/vision loss







### **Drugs in Development**

- Current drugs have limited efficacy with ~45% of patients being legally blind after 7 years.
- Other anti-fibrotic and anti-angiogenic agents currently in development with multiple targets
  - Platelet-Derived Growth Factor Receptor
  - Anti-ανβ1 integrin
  - Anti-ανβ3 integrin
  - Androgen Receptor Agonist
  - Angiopoietin 2 (ANGPT2) Inhibitor, Placental Growth Factor (PGF) Inhibitor, Vascular Endothelial Growth Factor A (VEGFA) Inhibitor
  - Sphingosine 1-Phosphate (S1P) Inhibitor
  - C5 Complement Inhibitor
  - Sodium Hydrogen Exchange Isoform-3 (NHE-3) Inhibitor
  - Connective Tissue Growth Factor (CTGF) Inhibitor





#### Fibrosis of the retina

- ► Fibrosis is a major factor leading to vision loss in a number of conditions including:
  - Treatment-resistant age related macular degeneration;
  - End stage diabetic retinopathy; and
  - Surgical repair of retinal detachment and trauma.
- Few if any current treatments
- ► Common causes including role of immune cells, tissue remodelling that is similar to fibrosis in other regions of the body







## Getting a fibrosis drug to the clinic and a deal

Stewart Roberts (chair)

Dr Brian Richardson

Dr John Westwick

Dr Robert Peach



#### Stuart Roberts, Chair

Stuart Roberts has been involved in the healthcare and biotechnology sector since the early 2002, initially as a sell-side analyst doing equities research in the sector in Australia, then, from the start of 2015, as an executive inside biotech companies, before he returned to equities research with the founding of NDF Research in June 2016.



#### Brian Richardson

Brian was most recently a member of The Leadership Team and The Global Head of The Musculoskeletal Disease Therapeutic Area at The Novartis Institutes for Biomedical Research having previously held several other senior positions during a 42 year career in the pharmaceutical industry.

Research conducted in Brian's laboratories has led to the discovery, development and introduction of several new therapies.



#### John Westwick

John has extensive experience in drug discovery in the Pharmaceutical Industry and as a Professor of Pharmacology. With over 14 years at Novartis Institutes for Biomedical Research, John was responsible for the build-up and leadership of all aspects of drug discovery and early development from target validation to the completion of proof of concepts in the respiratory area, which included severe asthma, Chronic Obstructive Pulmonary Disease (COPD), cystic fibrosis, pulmonary arterial hypertension, and pulmonary fibrosis.

John has had 13 positive proof of concepts in respiratory, which include a number of compounds and monoclonal antibodies which are now in phase III clinical trials.




#### Robert Peach

- Dr Peach has over 25 years of drug discovery and development experience in the pharmaceutical and biotechnology industry. In 2009 he co-founded Receptos, becoming Chief Scientific Officer and raising \$59M in venture capital and \$800M in an IPO and three subsequent follow-on offerings.
- ▶ In August 2015 Receptos was acquired by Celgene for \$7.8B. His extensive drug discovery and development experience in autoimmune and inflammatory diseases, and cancer has resulted in multiple drugs entering clinical trials and 3 registered drugs.



# **Drug development pathway**

- What does it take to get a fibrosis drug to the clinic?
- What does orphan drug designation mean?
- Which fibrosis indication?





## Fibrosis deals: IPF, NASH, Renal

**IPF** 



Sep-15 acquired by Roche \$105m + \$475m milestones phase I asset



Aug-15 acquired by BMS \$150m + \$1.25b milestones phase IIa asset

#### Galecto Biotech AB

Nov-14 acquired by BMS \$444m phase I asset

**NASH** 



May-15 acquired by BI \$40m + \$750m milestones phase I asset

### **NIMBUS**

**THERAPEUTICS** 

April-16 acquired by Gilead \$400m + \$800m milestones phase I asset



Sep-16 acquired by Allergan \$1.695B aquisition phase I & III assets

**RENAL** 



Sep-14 acquired by Shire \$75m + \$483m milestones phase I asset



August-14 acquired by Mallinckrodt \$5.6B





**i-bodies** – a new class of protein therapeutics to treat human disease

February 2017

Sam Cobb, CEO and Managing Director

AdAlta Limited (ASX:1AD)

s.cobb@adalta.com.au



### Sam Cobb

- ▶ Sam is the founding CEO of AdAlta and has over fifteen years' experience in business development and commercialisation of early stage scientific technologies.
- ▶ Prior to AdAlta, Sam was the Business Development Director at the Co-operative Research Centre for Diagnostics. Sam has also worked for the biotech start up companies Sensologix Inc and Nephrogenix Pty Ltd and at the University of Queensland's technology commercialisation companies, Uniquest Pty Ltd and IMBcom Pty Ltd.
- ▶ Sam has a Bachelor of Science, a Masters of Intellectual Property Law and has completed the Australian Institute of Company Directors course.

### **Disclaimer**

Investment in AdAlta is subject to investment risk, including possible loss of income and capital invested. AdAlta does not guarantee any particular rate of return or performance, nor do they guarantee the repayment of capital.

This presentation is not an offer or invitation for subscription or purchase of or a recommendation of securities. It does not take into account the investment objectives, financial situation and particular needs of the investor. Before making any investment in AdAlta, the investor or prospective investor should consider whether such an investment is appropriate to their particular investment needs, objectives and financial circumstances and consult an investment advisor if necessary.

This presentation may contain forward-looking statements regarding the potential of the Company's projects and interests and the development and therapeutic potential of the company's research and development. Any statement describing a goal, expectation, intention or belief of the company is a forward-looking statement and should be considered an at-risk statement. Such statements are subject to certain risks and uncertainties, particularly those inherent in the process of discovering, developing and commercialising drugs that are safe and effective for use as human therapeutics and the financing of such activities. There is no guarantee that the Company's research and development projects and interests (where applicable) will receive regulatory approvals or prove to be commercially successful in the future. Actual results of further research could differ from those projected or detailed in this presentation. As a result, you are cautioned not to rely on forward-looking statements. Consideration should be given to these and other risks concerning research and development programs referred to in this presentation.



# **Corporate and investment summary**

A drug discovery and development company focused on using its proprietary technology platform to generate a new class of protein therapeutics, known as i-bodies, for treating a wide range of human diseases

#### Investment highlights

- Initial focus on treating fibrosis high unmet medical need
- Advanced lead fibrosis drug candidate AD-114 with significant pre-clinical validation
- ► Fully funded for phase 1 development of lead fibrosis drug and i-body pipeline
- Early commercialisation potential
- Experienced team with strong track record of drug development and ability to deliver

Capital structure	
ASX code	1AD
Shares on issue*	101,037,617
Share price (31 January)	AU\$0.21
Market capitalisation	AU\$21m
Current cash	AU\$8.77m
Trading Range	AU\$0.31 to \$0.165

<sup>\* 50.9</sup>m shares escrowed for 6-24 months from listing


Major Shareholders	%
Yuuwa Capital LP	53.5
Platinum Asset Management	7.97
Citycastle Pty Ltd	5.26
La Trobe University	3.01
Robin Beaumont	1.82
Other shareholders	28.44
Total	100%



# Fibrosis: unmet medical need with multiple indications

- Developing i-bodies as improved therapies for the treatment of fibrosis
  - a condition that is prevalent in 45-50% of all diseases
- ► Fibrosis can occur in many tissues of the body as a result of inflammation or damage
  - it can result in scarring of vital organs causing irreparable damage and eventual organ failure
- AdAlta's initial focus is on lung fibrosis

Collectively fibrosis represents a large unmet clinical need



### Global market interest in fibrosis treatments


Recent transactions confirm that big pharma are actively acquiring fibrosis assets at an early stage – typically based on Phase I results

Date	Company	Target	Acquired by	Deal value (US\$)	Deal commentary
Sep-15	Adheron Therapeutics	SDP051	Roche	\$105M upfront, plus \$475M in milestones	SDP-51 at end of Phase I for IPF
Aug-15	Promedior	PRM-151	BMS	\$150m upfront + \$1.25B	Phase II IPF and myelofibrosis
Nov-14	Galecto Biotech AB	TD139	BMS	\$444M	Option to acquire at end of clinical POC (no later than 60 days following Ph 1b for IPF completion)
Aug-14	Intermune	Esbriet / Pirfenidone	Roche	\$8.3B	Approval in Europe / Japan, phase III in the US
Jun-13	MicroDose Therapeutx	MMI0100	Teva Pharmaceuticals	\$40M upfront \$125M milestones	MMI0100 was in pre-clinical development
Mar-12	Stromedix	STX100	Biogen Idec	\$75M upfront \$487.5M milestones	End of phase I for IPF
Jul-11	Amira / BMS	BMS-986020	BMS	\$325M upfront \$150M milestones	End of phase I for IPF

Source: Medtrack Pharma Intelligence, Informa (all IPF deals since 2011)



## **AD-114** development: key milestones



## **Expected news flow next 12 months**

Q3 & Q4 2016

- Commence manufacturing of material for toxicology testing with FujiFilm Diosynth Biotechnologies
- Additional AD-114 IPF fibrosis data
- ✓ Completion of evaluation of AD-114 with IPF clinicians Alfred Hospital
- ✓ Completion of AD-114 NASH animal study

H1 2017


- ✓ Orphan Drug Designation (US FDA)
- Hypertrophic scarring animal results for AD-114
- Manufactured material for toxicology testing available

H2 2017

- Eye fibrosis additional data, funded by NHMRC development grant
- Completion of other pre-clinical study animal models of AD-114
- ► AD-114 toxicology results



# AdAlta business model – strategy to create value



#### Market benchmarks

**Fibrosis** lead AD-114



Sep-15 acquired by Roche \$105m + \$475m milestones phase I asset



Aug-15 acquired by BMS \$150m + \$1.25b milestones phase IIa asset

#### Galecto Biotech AB

Nov-14 acquired by BMS \$444m phase I asset

Next gen antibodies



April-16 with Abbvie \$40m upfront + \$645m milestones & royalties



Dec -15 with Roche \$6.4m upfront + \$410m milestones & royalties



Nov-15 with Novo-Nordisk €9m upfront + €182m milestones & royalties)

**GPCRs** 



Acquired Feb-15 by Sosei \$400m Phase Ib asset + 7 preclinical leads



receptos

Acquired by Celgene July-15 \$8b Ph III, Ph II and GPCR platform



April-16 with Boehringer €8m payment for Ph1 GPCR nanobody (€125m milestones & royalties)



# Management and Board in place to deliver strategy



Sam Cobb: Founding CEO and Director

Extensive experience in raising equity, contract and grant funding

15 years of commercialisation and management experience



**Dr John Chiplin: Independent Director** 

CEO of investment Company NewStar Ventures

Managing Director of acquired antibody company

Arana Therapeutics



Dr Paul MacLeman: Chairman

Managing Director of a ASX listed IDT Australia Ltd

Founded biologics companies, experienced ASX listed executive



## Liddy McCall & Dr James Williams: Yuuwa Capital Directors

Founders and investment Directors of Yuuwa Capital

Founders of iCeutica Inc (acquired 2011) and Dimerix Limited



**Dr Robert Peach** 

Founder and CSO of Receptos Inc, acquired by Celgene Corporation in 2015 for US\$7.8bn

Deep experience in research and drug development



Directors of several Australian biotech and Agritech companies

Multiple FDA, CE Mark and TGA approvals



# **Scientific Advisory Board**

Internationally recognised with proven track record of drug development



David McGibney: pre-clinical and clinical advisor

20 years with Pfizer, including Head of European R&D

Ex Pfizer Ltd board member

Developed Viagra, and 10+ blockbuster drugs



## John Westwick: pulmonary drug discovery and development

Over 14 years experience at Novartis, head of respiratory drug discovery

Five product launches and 13 positive proof of concepts in respiratory, including a number of antibodies which are now in phase III.



## Brian Richardson: drug discovery and development expert

Ex-Sandoz and Novartis (40+ years), including Head of Pre-clinical Research

Over 60 original peer reviewed research papers



Dr Mick Foley, AdAlta CSO

Expert in phage display

NIH, NHMRC, ARC, Gates funding and over 70 scientific publications



# AdAlta investment summary

- Powerful proprietary technology platform to develop a pipeline of i-bodies for the treatment of a wide range of human diseases
- Initial focus on treating Idiopathic Pulmonary Fibrosis and other fibrotic diseases high unmet clinical need
- Advanced lead candidate with significant pre-clinical validation of AD-114 demonstrating anti-fibrotic and anti-inflammatory effects
- Early commercialisation opportunity
- Experienced management and Board to drive AD-114 development and secure technology platform partnerships and product licensing deals
- ▶ IPO August 2016 raised \$10M to meet major milestones: clinical trials of AD-114 in lung fibrosis and development of i-body pipeline





#### Thank-you

Sam Cobb, CEO and Managing Director

s.cobb@adalta.com.au

www.adalta.com.au